

AD-A110 458

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

BORSUK-ULAN THEOREMS FOR ARBITRARY S^1 ACTIONS AND APPLICATIONS. (U)

NOV 81 E R FADELL, S Y MUSSEINI


DAAG29-80-C-0041

NL

UNCLASSIFIED

MRC-TSR-2301

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178



LEVEL ~~IV~~

(2)

AD A110458

MA 758-

MRC Technical Summary Report #2301

BORSUK-ULAM THEOREMS FOR ARBITRARY
 S^1 ACTIONS AND APPLICATIONS

E. R. Fadell, S. Y. Hussein,
P. H. Rabinowitz

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

November 1981

Received June 1, 1981

DTIC FILE COPY

DTIC
ECTE
FEB 4 1982
H

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, D. C. 20550

221510 82 02 03 087

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

BORSUK-ULAM THEOREMS FOR ARBITRARY S^1 ACTIONS AND APPLICATIONS

E. R. Fadell, S. Y. Hussein, and P. H. Rabinowitz

Technical Summary Report #2301
November 1981

ABSTRACT

An S^1 version of the Borsuk-Ulam Theorem is proved for a situation where $\text{Fix } S^1$ may be nontrivial. The proof is accomplished with the aid of a new relative index theory. Applications are given to intersection theorems and the existence of multiple critical points is established for a class of functionals invariant under an S^1 symmetry.

AMS(MOS) Subject Classification: 55B25, 58E05

Key Words: Equivariant cohomology, index theory, Borsuk-Ulam theorem, intersection theorems, minimax, critical point, Hamiltonian system.

Work Unit No. 1 -Applied Analysis.



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041. This material is based upon work supported by the National Science Foundation under Grant No. MCS-8110556.

SIGNIFICANCE AND EXPLANATION

Minimax arguments from the calculus of variations serve as an important tool in establishing the existence of nonlinear vibrations of discrete mechanical systems as modelled by Hamilton's equations. In these arguments one obtains the solutions of the differential equations as critical points of an associated Lagrangian by minimaxing the Lagrangian over appropriate classes of sets. Intersection theorems such as are proved in this paper play a crucial role in this process. In addition to obtaining some intersection theorems we illustrate their use by proving an existence theorem for multiple critical points of a functional invariant under an S^1 symmetry group.

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

BORSUK-ULAM THEOREMS FOR ARBITRARY S^1 ACTIONS AND APPLICATIONS

E. R. Fadell, S. Y. Husseini and P. H. Rabinowitz

Introduction.

One of the variants of the Borsuk-Ulam Theorem states that if Ω is a bounded neighborhood of 0 in \mathbb{R}^n which is symmetric with respect to the origin and f is a continuous odd map of $\partial\Omega$ into a proper subspace of \mathbb{R}^n , then f has a zero on $\partial\Omega$ [1]. An extension of this result to an infinite dimensional setting for a class of Fredholm maps was carried out by Granas [2] and more quantitative versions of the result which provide lower bounds for a topological measure of the size of $f^{-1}(0) \cap \partial\Omega$ both in finite and infinite dimensions have been given by Holm and Spanier [3]. (See also [4]).

Our main goal in this paper is to obtain analogous results when Ω is invariant and f equivariant with respect to an S^1 rather than a \mathbb{Z}_2 action. For a class of such actions which are fixed point free on $\mathbb{R}^{2n} \setminus \{0\}$, available tools such as the index theory of [5] lead to an S^1 version of the Borsuk-Ulam Theorem by merely repeating the arguments of [3] or [4]. However when the action is not free, this approach fails. Nevertheless we will show how a relative index theory related to the cohomological index theory of [5] can be used to obtain a Borsuk-Ulam Theorem for a class of non-free S^1 actions. In §1 some properties of this index theory will be developed in a restricted setting. (A more systematic development will be carried out in a future paper). Some S^1 versions of the Borsuk-Ulam Theorem will be proved in §2. A special case is the following analogue of the \mathbb{Z}_2 result:

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041. This material is based upon work supported by the National Science Foundation under Grant No. MCS-8110556.

Theorem: Let S^1 act linearly on $\mathbb{R}^l \times \mathbb{R}^{2k}$ (i.e. via a group of unitary operators) so that $\text{Fix } S^1 = \mathbb{R}^l \times \{0\}$. Suppose that A is an annulus in $\mathbb{R}^l \times \mathbb{R}^{2k}$ and $f: A \rightarrow \mathbb{R}^l \times \mathbb{R}^{2k'}$, $k' < k$, is an equivariant map to a proper invariant subspace with $f|_{A^G} = \text{id}$, $A^G = (\text{Fix } S^1) \cap A$. If Ω is a closed, bounded invariant neighborhood of the origin with $\partial\Omega \subset A$, then $f^{-1}(0) \cap \partial\Omega \neq \emptyset$, i.e., f has zeroes on $\partial\Omega$.

The motivation for this paper was a study of the existence of periodic solutions of Hamiltonian systems of ordinary differential equations [6]. Minimax arguments from the calculus of variations furnish an important tool for treating such questions. In these arguments one obtains solutions of the differential equations as critical points of an associated functional by minimizing the functional over appropriate classes of sets. Intersection theorems play a crucial role in getting estimates, in particular lower bounds, on minimax values and in showing they are indeed critical values of the functional. In §3 the Borsuk-Ulam results will be used to obtain some new intersection theorems. To illustrate their use an abstract critical point theorem (see Theorem 3.14 below) in an S^1 setting related to a \mathbb{Z}_2 result from [7] will be proved.

§1. A Relative Index

Throughout this section $G = S^1$, the group of complex numbers of norm 1. Suppose X is a paracompact G -space, i.e. $G (= S^1)$ acts on X . As usual, the fixed point set under this action is given by

$$X^G = \{x \in X : gx = x \text{ for all } g \in G\}.$$

Also, for $x \in X$, the isotropy subgroup at $x \in X$ is given by

$$G_x = \{g \in G : gx = x\}.$$

Note, that because $G = S^1$, G_x is a finite cyclic group when $x \notin X^G$.

Following [5], let $\eta = (E_G, p, B_G)$ denote the universal G -bundle, e.g.

$E_G = S^\infty$, $B_G = CP^\infty$, see [5]. Set

$$E_G(X, X^G) = E_G \times_G (X, X^G) = (E_G \times_G X, B_G \times X^G)$$

where G acts freely on $E_G \times X$ and $E_G \times X^G$ by the action $g(e, x) = (ge, gx)$ and $E \times_G X$, $B_G \times X^G$ are the resulting orbit spaces. The projection $E_G \times X \rightarrow E_G$ induces a fiber bundle pair

$$(q, q_0) : E_G(X, X^G) \rightarrow B_G$$

with fiber (X, X^G) . $q : E \times_G X \rightarrow B_G$ is, in fact, the classifying map of the principal G -bundle, $\eta : E_G \times X \rightarrow E_G \times_G X$ and $q_0 : B_G \times X^G \rightarrow B_G$ is simply projection on B . Now, using Alexander-Spanier cohomology H^* with rational coefficients, we set

$$H_G^*(X, X^G) = H^*(E_G(X, X^G))$$

and

$$H_G^*(X) = H^*(E_G(X)) = H^*(E_G \times_G X).$$

The relative cup product

$$H_G^*(X) \times H_G^*(X, X^G) \rightarrow H_G^*(X, X^G)$$

then gives $H_G^*(X, X^G)$ a module structure over the ring $\Lambda = H^*(B_G) =$

$H_G^*(\text{point})$ with left action given by

$$\lambda \cdot z = q^*(\lambda) \cup z, \quad \lambda \in \Lambda, \quad z \in H_G^*(X, X^G).$$

We recall also the fact that $\Lambda = Q[\alpha]$, where $Q[\alpha]$ is the polynomial algebra on a single generator $\alpha \in H^2(B_G)$.

Definition (1.1) The relative α -index $\text{Ind}_\alpha(X, X^G)$ of the pair (X, X^G) is the smallest integer s such that α^s annihilates $H_G^*(X, X^G)$, i.e.

$$\text{Ind}_\alpha(X, X^G) = \min_s \{ \alpha^s \cdot x = 0 \text{ for all } x \in H_G^*(X, X^G) \}.$$

Proposition (1.2). When $X^G = \emptyset$, the relative index coincides with the α -index in [5], i.e.

$$\text{Ind}_\alpha(X, \emptyset) = \text{Index}_\alpha^* X$$

and hence in this special situation all the basic properties of Index_α^* in [5] hold for $\text{Ind}_\alpha(X, \emptyset)$.

Proof. When $X^G = \emptyset$, $H_G^*(X, X^G) = H_G^*(X)$ and we have a unit $1 \in H_G^0(X)$. Then, since

$$\alpha^s \cdot 1 = 0 \iff \alpha^s = 0$$

we have the desired result.

Remark (1.3). The basic properties of the relative index $\text{Ind}_\alpha(X, X^G)$ in general require some modification and will be taken up in a future work. They will be studied in the category of pairs (X, A) where X is a G -space, A an invariant subspace and G an arbitrary compact Lie group. The sequel will not require these modified properties. However, the following one is of interest since in [5], $X^G \neq \emptyset$ always implies $\text{Index}_\alpha^* X = \infty$.

Proposition (1.4). If X is a finite dimensional separable metric space, then

$$\text{Ind}_\alpha(X, X^G) < \infty.$$

Proof. The natural projection $E \times X \rightarrow X$ induces a map

$$\varphi : E_G(X, X^G) \rightarrow (X/G, X^G)$$

which induces isomorphisms [15]

$$\varphi^* : H^*(X/G, X^G) \xrightarrow{\cong} H_G^*(X, X^G)$$

(because $G = S^1$ and coefficients are rational). When X is a finite dimensional separable metric space, so is X/G and hence $H_G^n(X, \mathbb{Q})$ vanishes for n sufficiently large.

We now proceed to calculate an important example. Suppose that $G = S^1$ acts linearly on $\mathbb{R}^n = \mathbb{R} \times \cdots \times \mathbb{R}$, the n -fold cartesian product of the real line. Introduce a G -equivariant inner-product on \mathbb{R}^n (by the "unitary trick") and assume that $G = S^1$ is a subgroup of $SO(n, \mathbb{R})$, the group of rotations of \mathbb{R}^n . The fixed-point set $(\mathbb{R}^n)^G = \{x \in \mathbb{R}^n \mid \sigma x = x, \text{ for all } \sigma \in G\}$ is a linear subspace \mathbb{R}^l of dimension, say l . Then clearly the orthogonal complement $(\mathbb{R}^l)^\perp$ is G -invariant and of even-dimension, say $2n'$. Thus we obtain the G -equivariant decomposition

$$\mathbb{R}^n = \mathbb{R}^l \times \mathbb{R}^{2n'}$$

with $\mathbb{R}^l \times \{0\} = (\mathbb{R}^n)^G$. Note that if $x \notin \mathbb{R}^l \times \{0\}$ then the isotropy subgroup $G_x \subset G$ is a finite cyclic subgroup of G . We wish to compute

$\text{Ind}_G(S^{n-1}, (S^{n-1})^G)$, where S^{n-1} is the unit sphere in \mathbb{R}^n ,

$$(S^{n-1})^G = S^{n-1} \cap (\mathbb{R}^l \times \{0\}) = S^{l-1}$$

and S^{l-1} is the unit sphere in $\mathbb{R}^l \times \{0\}$.

Proposition (1.5). $\text{Ind}_G(S^{n-1}, S^{l-1}) = n'$, where $n = l + 2n'$.

Note that if $l = 0$, then Proposition (1.5) is just Proposition (7.4) of [5]. We shall prove Proposition (1.5) in stages. First, let us consider the fibration

$$q : E_G(S^{n-1}) \rightarrow B_G$$

with fiber S^{n-1} .

Lemma (1.6): There exists an isomorphism of $H^*(B_G)$ -modules

$$\varphi : H^*(B_G) \otimes_{\mathbb{Q}} H^*(S^{n-1}) \rightarrow H_G^*(S^{n-1})$$

given by $\varphi(u \otimes x) = q^*(u) \cup \theta(x)$, where $\theta : H^*(S^{n-1}) \rightarrow H_G^*(S^{n-1})$ is a cohomology extension of the fiber. Let $\bar{\gamma}$ denote a generator of $H^{n-1}(S^{n-1})$

and $\gamma = \varphi(1 \otimes \bar{\gamma})$. Then, for certain integers r and t and $c_1 \in \mathbb{Q}$,

$$\gamma^2 = \gamma(c_1 \alpha^r + c_2 \alpha^t \otimes \bar{\gamma}).$$

Proof. This is an immediate consequence of the Leray-Hirsch Theorem [15] as soon as we establish the triviality of the (rational) spectral sequence of the fibration $S^{n-1} \rightarrow E \times_G S^{n-1} \rightarrow B_G$. The differential operators

$d_r^{p,q} : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$ vanish for $2 \leq r \leq n-1$ and in fact the only possible non-trivial differential operator occurs when $r = n$. Consider the diagram

$$\begin{array}{ccc} E_G(S^{l-1}) & \xrightarrow{\quad} & E_G(S^{n-1}) \\ & \searrow q' & \swarrow q \\ & B_G & \end{array}$$

Since $E_G(S^{l-1}) = S^{l-1} \times B_G$, $(q')^*$ and hence q^* is injective. If the differential operators

$$d_n^{p,n-1} : E_n^{p,n-1} \rightarrow E_n^{p+n,0}$$

were not trivial, then the edge homomorphism diagram

$$\begin{array}{ccc} H^{p+n}(B_G) = E_n^{p+n,0} & \xrightarrow{\quad} & E_{n+1}^{p+n,0} = E_{\infty}^{p+n,0} \\ & \searrow q^* & \swarrow \\ & H_G^{p+n}(S^{n-1}) & \end{array}$$

would produce a non-trivial kernel for q^* . Thus all d_r 's are trivial.

Remark (1.7). If $(n-1)$ is odd, then $\gamma^2 = 0$ above.

We consider now the exact sequence of the pair

$$E_G(S^{n-1}, S^{l-1})$$

$$\cdots \rightarrow H_G^q(S^{n-1}, S^{l-1}) \xrightarrow{j^*} H_G^q(S^{n-1}) \xrightarrow{i^*} H_G^q(S^{l-1}) \rightarrow \cdots$$

Lemma (1.8). i^* is a monomorphism.

Proof. First observe that j^* and i^* are $H^*(B_G)$ -homomorphisms. Suppose $i^*(x) = 0$. Then, for some element $u \in H_G^*(S^{n-1}, S^{l-1})$ $j^*(u) = x$. But since $H_G^r(S^{n-1}, S^{l-1})$ vanishes for r large, $j^*(\alpha^r(u)) = 0 = \alpha^r(j^*(u)) = 0$. But $H^*(B_G)$ operates freely on $H_G^*(S^{n-1})$ and hence $j^*(u) = 0 = x$.

Lemma (1.9). $H_G^*(S^{n-1}, S^{l-1})$ is generated over $H^*(B_G)$ by a single element β with $\alpha^{n'}\beta = 0$, i.e.

$$H_G^*(S^{n-1}, S^{l-1}) = \{\beta, \alpha\beta, \dots, \alpha^{n'-1}\beta \mid \alpha^{n'}\beta = 0\}.$$

Proof. Consider the short exact sequence of $H^*(B_G)$ -modules

$$0 \rightarrow H_G^q(S^{n-1}) \xrightarrow{i^*} H_G^q(S^{l-1}) \xrightarrow{\delta^*} H_G^{q+1}(S^{n-1}, S^{l-1}) \rightarrow 0$$

and $H^*(B_G)$ -isomorphisms

$$\varphi : H^*(B_G) \otimes H^*(S^{n-1}) \rightarrow H_G^*(S^{n-1})$$

$$\psi : H^*(B_G) \otimes H^*(S^{l-1}) \rightarrow H_G^*(S^{l-1}).$$

Let $\bar{\gamma}$ and $\bar{\beta}$ denote the fundamental class of S^{n-1} and S^{l-1} , respectively, and $\gamma = \varphi(1 \otimes \bar{\gamma})$, $\beta = \psi(1 \otimes \bar{\beta})$. Then, we have two cases:

$$i^*(\gamma) = \psi(c\alpha^{n'} \otimes \beta + d\alpha^{\frac{n-1}{2}}) \text{ if } n-1 \text{ is even}$$

and

$$i^*(\gamma) = \psi(c\alpha^{n'} \otimes \beta) \text{ if } n-1 \text{ is odd.}$$

In either case $\psi(\alpha^{n'} \otimes \beta)$ is in the image of i^* while the elements

$\psi(\alpha^j \otimes \beta)$, $j < n'$, are not. This gives the desired result.

Proof of Proposition (1.5). Apply the previous lemma.

Proposition (1.5) is the key to the following Theorem. (cf. Proposition (3.9) of [5].)

Theorem (1.10): Suppose that G acts on $R^n = R^l \times R^{2n'}$ as described above, and let Ω be a closed, bounded invariant neighborhood of 0 in R^n . Then

$$\text{Ind}_G(\partial\Omega, (\partial\Omega)^G) > n'.$$

Theorem (1.10) is an immediate consequence of the following form of the Piercing Property [5]. Notice the conclusion is $>$ and not equality.

Proposition (1.11) (Piercing Property). Let X denote a paracompact G -space and $X \times I$ the corresponding G -space with action $g(x,t) = (gx,t)$, $g \in G$, $x \in X$, $t \in I$. Let $A_0 = X \times \{0\}$, $A_1 = X \times \{1\}$. Suppose

$$X \times I = B_0 \cup B_1, \quad A_0 \subset B_0, \quad A_1 \subset B_1$$

where B_0 and B_1 are closed invariant subsets. Then, if $C = B_0 \cap B_1$,

$$\text{Ind}_G(C, C^G) > \text{Ind}_G(X, X^G).$$

Proof. The proof is essentially the same as in [5]. Let $B = B_0 \cup B_1$. Then the inclusion maps

$$\begin{array}{ccc} A_0 & \xrightarrow{k_0} & B_0 \\ j_0 \searrow & & \nearrow i_0 \\ & B & \end{array} \quad \begin{array}{ccc} A_1 & \xrightarrow{k_1} & B_1 \\ j_1 \searrow & & \nearrow i_1 \\ & B & \end{array} \quad \begin{array}{l} \ell_0 : C \subset B_0 \\ \ell_1 : C \subset B_1 \end{array}$$

induce $H^*(B_G)$ -homomorphisms, $k_0^* : H_G^*(B_0, B_0^G) \rightarrow H_G^*(A_0, A_0^G)$, j_0^* , i_0^* , etc.

Consider the Mayer-Vietoris sequence

$$\dots \rightarrow H_G^q(B, B^G) \xrightarrow{\zeta} H_G^q(B_0, B_0^G) \oplus H_G^q(B_1, B_1^G) \xrightarrow{\eta} H_G^q(C, C^G) \rightarrow$$

where $\zeta = (i_0^*, -i_1^*)$, $\eta = \ell_0^* + \ell_1^*$. If $\ell_0^*(x) = 0$, then $\eta(x, 0) = 0$ and

$i_0^*(y) = x$, $i_1^*(y) = 0$ for some y . But then $j_1^*(y) = 0$. Since $j_1 : (A_1, A_1^G) \rightarrow (B, B^G)$ is an equivariant homotopy equivalence,

$j_1^* : H_G^*(B, B^G) \rightarrow H_G^*(A_1, A_1^G)$ is an isomorphism and hence $y = 0$. Thus $x = 0$ and ℓ_0^* is a monomorphism. Now, let $\gamma : B \rightarrow A_0$ denote projection (which is equivariant) and $\gamma_C : C \rightarrow A_0$, $\gamma_0 : B_0 \rightarrow A_0$ the corresponding restrictions. Notice that we do not conclude that γ_C implies $\text{Ind}_G(C, C^G) <$

$\text{Ind}_G(A_0, A_0^G)$. However, in the diagram

$$\begin{array}{ccc}
 H_G^q(B_0, B_0^G) & \xrightarrow{\ell_0^*} & H_G^q(C, C^G) \\
 \gamma_0^* \swarrow & & \nearrow \gamma_C^* \\
 & H_G^q(A_0, A_0^G) &
 \end{array}$$

γ_0^* is also a monomorphism which forces γ_C^* to be a monomorphism of $H^*(B_G)$ -modules. Thus if $\alpha^s \cdot \gamma_C^*(c) = 0$, then $\alpha^s c = 0$ and since $(A_0, A_0^G) \equiv (X, X^G)$,
 $\text{Ind}_\alpha(C, C^G) > \text{Ind}_\alpha(X, X^G)$.

Proof of (1.10). Let A denote an annulus in \mathbb{R}^n centered at the origin with $\partial\Omega \subset \text{int } A$. Then, $A \approx B_0 \cup B_1$ where $B_0 = A \cap \Omega$, $B_1 = A \cap (\mathbb{R}^n \setminus \text{int } \Omega)$ and $\partial\Omega = B_0 \cap B_1$. Thus, applying (1.11) we have

$$\text{Ind}_\alpha(\partial\Omega, (\partial\Omega)^G) > n'.$$

We next consider special forms of the monotonicity and additivity properties which will be useful in the next section.

Proposition (1.12). Let X and Y denote paracompact G -spaces and $f: X \rightarrow Y$ an equivariant map. If

$$f^* : H_G^*(Y, Y^G) \rightarrow H_G^*(X, X^G)$$

is surjective, then $\text{Ind}_\alpha(X, X^G) \leq \text{Ind}_\alpha(Y, Y^G)$.

Proof. Suppose $\text{Ind}_\alpha(Y, Y^G) = s$. Then, $f^*(\alpha^s y) = \alpha^s f^*(y) = 0$. Since every element of $H_G^*(X, X^G)$ is of the form $f^*(y)$, α^s annihilates $H_G^*(X, X^G)$ and the result follows.

Proposition (1.13). Let X denote a paracompact G -space and A and B paracompact subsets whose interiors cover X . Suppose further that $B^G = \emptyset$ (so that all the fixed points of X are in A). Then,

$$\text{Ind}_\alpha(X, X^G) \leq \text{Ind}_\alpha(A, A^G) + \text{Ind}_\alpha(B, \emptyset).$$

Proof. Suppose $\text{Ind}_\alpha(A, A^G) = a$, $\text{Ind}_\alpha(B, \emptyset) = b$ where both a and b are finite. Consider the diagram

$$\begin{array}{ccc} E_G(A) & \xrightarrow{i_1} & E_G(X) \\ & \nearrow i_2 & \xrightarrow{f} B_G \\ E_G(B) & & \end{array}$$

where i_1 and i_2 are induced by inclusions and f is the classifying map for $E \times G \rightarrow E_G(X)$. Then, if $\beta = i_1^* f^*(\alpha)$ and $\gamma = i_2^* f^*(\alpha)$ we have β^a annihilates $H_G^*(A, A^G)$ and $\gamma^b = 0$. Take any element

$$x \in H_G^D(X, X^G)$$

and consider the exact sequences ($X^G = A^G$)

$$\cdots \rightarrow H_G^r(X, A) \xrightarrow{j_1^*} H_G^r(X, X^G) \rightarrow H_G^r(A, A^G) \rightarrow$$

$$\cdots \rightarrow H_G^s(X, B) \xrightarrow{j_2^*} H_G^s(X) \rightarrow H_G^s(B) \rightarrow .$$

Then, $f^*(\alpha^a) \cup x$ and $f^*(\alpha^b)$ pull back under j_1^* and j_2^* , respectively.

Then the following diagram

$$\begin{array}{ccc} H_G^*(X, A) \oplus H_G^*(X, B) & \xrightarrow{U} & H_G^*(X, X) \\ \downarrow & & \downarrow \\ H_G^*(X, X^G) \oplus H_G^*(X) & \xrightarrow{U} & H_G^*(X, X^G) \end{array}$$

shows that $f^*(\alpha^{a+b}) \cup x = 0$ and $\text{Ind}_\alpha(X, X^G) \leq a+b$.

Remark (1.14). (a) If $G = S^0 = \{\pm 1\}$, then using cohomology with coefficients in \mathbb{Z}_2 instead of \mathbb{Q} , one proves in a similar fashion to the above that

$$\text{Ind}_G(\partial\Omega, (\partial\Omega)^G) > n - l = 2n',$$

where $R^n = R^l \times R^{2n'}$.

(b) If $G = S^3$, the unit quaternions and $G_x \subset G$ is finite for $x \notin R^l \times \{0\}$, then one can prove, using rational cohomology, that

$$\text{Ind}_G(\partial\Omega, (\partial\Omega)^G) > n',$$

where $R^n = R^l \times R^{4n'}$. The general situation is much more complicated.

(c) If G is a finite cyclic group of prime order, then using cohomology with coefficients in \mathbb{Z}_p , one can prove that

$$\text{Ind}_G(\partial\Omega, (\partial\Omega)^G) > n',$$

where $R^n = R^l \times R^{2n'}$. If G does not have prime order, the problem is more complicated.

We close this section with the following result which was not included in [5] and will also prove useful.

Proposition (1.15). When X is compact and $X^G = \emptyset$, then

$$\text{Ind}_G(X, \emptyset) = \text{Index}_G^* X < \infty.$$

Proof. For $x \in X$, the orbit $Gx = G/G_x$, where G_x is the isotropy group at x . G_x is a finite cyclic group and $\text{Index}_G^*(Gx) = 1$ ([5], Proposition 6.12). A suitable neighborhood $N(Gx)$ of this orbit also has

$\text{Index}_G^* N(Gx) = 1$. Since X is compact, a finite number of such neighborhoods cover X and subadditivity ([5], Proposition 6.6) completes the proof.

2. Zeros of Equivariant Maps.

Suppose that $G (=S^1)$ acts on $\mathbb{R}^a = \mathbb{R}^l \times \mathbb{R}^{2a'}$ and $\mathbb{R}^b = \mathbb{R}^l \times \mathbb{R}^{2b'}$ as a group of rotations in such a way that (as before)

$$(\mathbb{R}^a)^G = \mathbb{R}^l \times \{0\}, \quad (\mathbb{R}^b)^G = \mathbb{R}^l \times \{0\}.$$

Suppose we have an equivariant map

$$f : S^{a-1} \rightarrow \mathbb{R}^b \setminus \{0\}$$

such that the restriction $f_0 : S^{l-1} \rightarrow \mathbb{R}^l \setminus \{0\}$ has the property that

$$f_0^* : H^{l-1}(\mathbb{R}^l \setminus \{0\}) \rightarrow H^{l-1}(S^{l-1})$$

is an isomorphism. Then, in the following diagram

$$\begin{array}{ccc} H_G^{l-1}(\mathbb{R}^l \setminus \{0\}) & \xrightarrow{\delta} & H_G^l(\mathbb{R}^b \setminus \{0\}, \mathbb{R}^l \setminus \{0\}) \\ (f_0)_G^* \downarrow & & \downarrow f_G^* \\ H_G^{l-1}(S^{l-1}) & \xrightarrow{\delta} & H_G^l(S^{a-1}, S^{l-1}) \end{array}$$

$(f_0)_G^*$ is an isomorphism and f_G^* takes the generator (over $H^*(B_G)$) to a generator and f_G^* is surjective. Applying the monotonicity property (1.12) we have

$$a' = \text{Ind}_G(S^{a-1}, S^{l-1}) \leq \text{Ind}_G(\mathbb{R}^b \setminus \{0\}, \mathbb{R}^l \setminus \{0\}) = b'$$

Thus, we have the following result.

Proposition (2.1). Suppose $f : S^{a-1} \rightarrow \mathbb{R}^b$ is an equivariant map, where \mathbb{R}^a and \mathbb{R}^b are as above. If

$$f_0^* : H^{l-1}(\mathbb{R}^l \setminus \{0\}) \rightarrow H^{l-1}(S^{l-1})$$

is an isomorphism, then $f^{-1}(0) \neq \emptyset$, i.e. f possesses at least one zero.

We will now extend this result to maps $f : \partial\Omega \rightarrow \mathbb{R}^b$ where Ω is a closed, bounded invariant subset of \mathbb{R}^a . Let A denote an annulus in \mathbb{R}^a (as usual, the region between two concentric spheres centered at the origin) and let Ω denote a closed, bounded invariant neighborhood of the origin with

$\partial\Omega \subset A$, where $\partial\Omega$ is the boundary relative to \mathbb{R}^a . We assume once and for all that

$$f : A \rightarrow \mathbb{R}^b$$

is a given equivariant map.

Proposition (2.2). Suppose $f(A^G) \subset \mathbb{R}^b \setminus \{0\}$ and

$$(f|A^G)^* : H^*(\mathbb{R}^b \setminus \{0\}) \rightarrow H^*(A^G)$$

is an isomorphism. Then, if $a > b$, $Z = f^{-1}(0) \cap \partial\Omega \neq \emptyset$.

Proof. Suppose $f|_{\partial\Omega} : \partial\Omega \rightarrow \mathbb{R}^b \setminus \{0\}$ and consider the diagram

$$\begin{array}{ccccc}
 & H_G^{l-1}(\mathbb{R}^b \setminus \{0\}) & \xrightarrow{\delta} & H_G^l(\mathbb{R}^b \setminus \{0\}, \mathbb{R}^b \setminus \{0\}) & \\
 & \downarrow (f|(\partial\Omega)^G)^* & & \downarrow (f|_{\partial\Omega})^* & \\
 (f|A^G)^* & H_G^{l-1}((\partial\Omega)^G) & \xrightarrow{\delta} & H_G^l(\partial\Omega, (\partial\Omega)^G) & \\
 & \uparrow i^* & & \uparrow j & \\
 & H_G^{l-1}(A^G) & \xrightarrow{\delta} & H_G^l(A, A^G) &
 \end{array}$$

The G-space A is equivariantly homotopic to S^{a-1} , and $\mathbb{R}^b \setminus \{0\}$ to S^{b-1} .

Choose

$$\beta_1 \in H_G^{l-1}(\mathbb{R}^b \setminus \{0\}), \quad \beta_2 = (f|A^G)^*(\beta_1) \in H_G^{l-1}(A^G)$$

so that $\beta_1 = \delta\tilde{\beta}_1$ and $\beta_2 = \delta\tilde{\beta}_2$ are generators over $H^*(B_G)$ of

$H_G^*(\mathbb{R}^b \setminus \{0\}, \mathbb{R}^b \setminus \{0\})$ and $H_G^*(A, A^G)$, respectively. Then, if $\tilde{\beta}_3 = i^*(\tilde{\beta}_2) =$

$(f|(\partial\Omega)^G)^*(\tilde{\beta}_1)$, $\beta_3 = \delta\tilde{\beta}_3$ generates a subalgebra isomorphic by j^* to

$H_G^l(A, A^G)$. Since $\beta_3 = (f|_{\partial\Omega})^*(\beta_1)$ we have $\alpha^{b'} \cdot \beta_3 = 0$. On the other

hand, $\alpha^{a'} \cdot \beta_3 = 0$ with a' minimal. Thus, $a' \leq b'$ and $a \leq b$. Thus,

$a > b$ implies $Z = f^{-1}(0) \cap \partial\Omega \neq \emptyset$.

We now proceed to determine the index of the zero set Z .

Theorem (2.3). Under the hypotheses of (2.2), if $Z = f^{-1}(0) \cap \partial\Omega$,

$$\text{Ind}_G Z \geq a' - b'.$$

Proof. We know $Z \neq \emptyset$ by Proposition (2.2), although this fact is not necessary for the argument. Suppose $s = \text{Ind}_\alpha Z < a' - b'$. Then, Z is a closed subset of $\partial\Omega$ and hence of $\partial\Omega \setminus (\partial\Omega)^G$. Since $Z^G = \emptyset$, $\text{Index}_\alpha^* Z = \text{Ind } Z$, where Index_α^* is the index theory in [5]. Thus, since Index_α^* is continuous, we have a closed neighborhood N of Z (in $\partial\Omega$) such that $N^G = \emptyset$, $\text{Ind}_\alpha N = s$ and a closed set $M \subset \partial\Omega$ such that $\text{int} M \cup \text{int} N = \partial\Omega$, $M \cap Z = \emptyset$ and $M^G = (\partial\Omega)^G$. We know that $H_G^*(\partial\Omega, (\partial\Omega)^G)$ contains an $H^*(B_G)$ -submodule $\{\beta, \alpha\beta, \alpha^2\beta, \dots, \alpha^{a'-1}\beta : \alpha^{a'}\beta = 0\}$. The inclusion map

$$j : (M, M^G) \rightarrow (\partial\Omega, (\partial\Omega)^G)$$

induces

$$j^* : H^*(\partial\Omega, (\partial\Omega)^G) \rightarrow H^*(M, M^G)$$

and we set $\beta_M = j^*(\beta)$. Let t denote the smallest integer such that

$\alpha^t \beta_M = 0$. The diagram

$$\begin{array}{ccc} H_G^{l-1}(\mathbb{R}^l \setminus \{0\}) & \xrightarrow{\delta} & H^l(\mathbb{R}^b \setminus \{0\}, \mathbb{R}^l \setminus \{0\}) \\ (f|M^G)^* \downarrow & & \downarrow (f|M)^* \\ H_G^{l-1}(M^G) & \xrightarrow{\delta} & H^l(M, M^G) \\ \uparrow \text{id} & & \uparrow \\ H_G^{l-1}((\partial\Omega)^G) & \xrightarrow{\delta} & H^l(\partial\Omega, (\partial\Omega)^G) \end{array}$$

tells us that $t \leq b'$. On the other hand using the argument for the additivity property (1.13) and the diagram

$$\begin{array}{ccc} H^*(\partial\Omega, N) \oplus H^*(\partial\Omega, M) & \rightarrow & H^*(\partial\Omega, \partial\Omega) \\ \downarrow & & \downarrow \\ H^*(\partial\Omega) \oplus H^*(\partial\Omega, (\partial\Omega)^G) & \rightarrow & H^*(\partial\Omega, (\partial\Omega)^G) \\ \downarrow & & \downarrow \\ H^*(N) \oplus H^*(M, M^G) & \nearrow & \end{array}$$

we see that $\alpha^{s+t}\beta = 0$ so that

$$a' < s+t < (a'-b') + b' = a'$$

which is a contradiction. Thus, $\text{Ind}_\alpha Z > a' - b'$.

Corollary (2.4). If $f: A \rightarrow K^b$ is the identity map on A^G and Z is as above, then

$$\text{Ind}_\alpha Z > a' - b'.$$

3. Applications.

In this section some applications of Theorem 2.3 will be given. First some finite and infinite dimensional intersection theorems will be obtained. Then these results will be used to prove an abstract critical point theorem in an S^1 setting. In the applications below we will be dealing with situations in which $X \cap \text{Fix } S^1 = \emptyset$. Hence $X^G = \emptyset$ and $\text{ind}_\alpha(X, X^G) = \text{Ind}_\alpha(X, \emptyset) = \text{Index}_\alpha^* X$ ([5]). As a convenience we list the properties of Ind_α from [5] (and (1.14)) we will require below, and at the same time suppress the α . We also let $C^k(X, Y)$ denote the set of k times continuously Frechet differentiable mappings from X to Y .

Lemma 3.1: Let E be a Hilbert space and let S^1 act on E . Let \mathcal{E} denote the family of equivariant subsets of $E \setminus \text{Fix } S^1$. Then $\text{Ind}: \mathcal{E} \rightarrow \mathbb{N} \cup \{\infty\}$ possesses the following properties: For $X, Y \in \mathcal{E}$,

- 1°. If $h \in C(X, Y)$ is equivariant, $\text{Ind } X \leq \text{Ind } Y$.
- 2°. $\text{Ind}(X \cup Y) \leq \text{Ind } X + \text{Ind } Y$.
- 3°. If X is compact, then $\text{Ind } X < \infty$ and there exists a $\delta > 0$ such that $\text{Ind } X = \text{Ind } N_\delta(X)$ where $N_\delta(X) = \{x \in E \mid \|x - X\| < \delta\}$.

Our first intersection result is in a Euclidean space setting. Let B_R^j denote the closed ball of radius R centered about 0 in $\mathbb{R}^l \times \mathbb{R}^{2j'}$ where $j = l + 2j'$. Whenever $j' < k'$, we will consider $\mathbb{R}^{2j'}$ as a subspace of $\mathbb{R}^{2k'}$ via $\mathbb{R}^{2j'} = \mathbb{R}^{2j'} \times \{0\} \subset \mathbb{R}^{2j'} \times \mathbb{R}^{2(k'-j')} = \mathbb{R}^{2k'}$.

Let S^1 act on $\mathbb{R}^l \times \mathbb{R}^{2k'}$ via a group of unitary operators such that $\text{Fix } S^1 = \mathbb{R}^l \times \{0\}$ and $\{0\} \times \mathbb{R}^{2j'}$ is an invariant subspace of

$\{0\} \times \mathbb{R}^{2k'}$ for $j' < k'$. Suppose $h \in C(B_R^k, \mathbb{R}^l \times \mathbb{R}^{2m'})$ where $m' > k'$, $h(x) = x$ on $(\mathbb{R}^l \times \{0\}) \cap B_R^k$, and h is equivariant. Let $\rho < R$ and consider $h^{-1}(B_\rho^m)$. This is an invariant neighborhood of 0 in $\mathbb{R}^l \times \mathbb{R}^{2k'}$. Let Ω denote the component of $h^{-1}(B_\rho^m)$ which contains 0. Then $\partial\Omega$ is an

invariant set. For $j' < k'$, let $P_{l+2j'}$ denote the orthogonal projector of $\mathbb{R}^l \times \mathbb{R}^{2m'}$ onto $\mathbb{R}^l \times \mathbb{R}^{2j'}$ and consider $f \equiv P_{l+2j'} h$. Then $f \in C(\partial\Omega, \mathbb{R}^l \times \mathbb{R}^{2j'})$, $f(x) = x$ for $x \in (\mathbb{R}^l \times \{0\}) \cap \Omega$, and f is equivariant. Thus as an immediate consequence of Theorem 2.3 we have

Proposition 3.2: $\text{Ind}(f^{-1}(0) \cap \partial\Omega) \geq k' - j'$.

With a bit more structure on h we have

Proposition 3.3: If $k \in C(B_R^k, \mathbb{R}^l \times \mathbb{R}^{2m'})$ where $m' \geq k'$, $h(x) = x$ for $x \in (\mathbb{R}^l \times \{0\}) \cup \partial B_R^k$, and h is equivariant, then for any $\rho < R$ and $j' < k'$,

$$\text{Ind}(h(B_R^k) \cap \partial B_\rho^m \cap (\{0\} \times \mathbb{R}^{2(m'-j')})) \geq$$

(3.4)

$$\text{Ind}(\{x \in B_R^k \mid h(x) \in \partial B_\rho^m \cap (\{0\} \times \mathbb{R}^{2(m'-j')})\}) \geq k' - j'.$$

Proof: We need only observe that $h(x) = x$ on ∂B_ρ^m implies Ω is strictly interior to B_ρ^m . Hence if $x \in \partial\Omega$, $h(x) \in \partial B_\rho^m$ via the maximality of Ω . Now (3.4) follows from 1° of Lemma 3.1 and Proposition 3.2.

Next we will prove two infinite dimensional extensions of Proposition 3.2. For what follows let E be a separable infinite dimensional Hilbert space and suppose S^1 acts on E , the action being given by a group G of unitary operators on E . Further assume $\text{Fix } G \equiv E_0$ is a finite dimensional subspace of E with $\dim E_0 = l$ and we can choose an orthogonal basis $\{v_j\}$ in E such that $E_0 = \text{span}\{v_1, \dots, v_l\}$, $E_m = \text{span}\{v_1, \dots, v_{l+2m}\}$, E_m is an invariant subspace of E , and $E = \overline{\bigcup_m E_m}$. For brevity when the above are satisfied, we say E, S^1 satisfy (*).

As an example, consider $W^{1,2}(S^1)$ the Hilbert space of 2π periodic functions under the norm

$$\|q\|^2 = \int_0^{2\pi} \left(\left| \frac{dq}{dt} \right|^2 + |q|^2 \right) dt.$$

There is a natural S^1 action on $W^{1,2}(S^1)$. Indeed for $q \in E$ and $\theta \in [0, 2\pi)$ let $T_\theta q(t) = q(t+\theta)$ and $G = \{T_\theta | \theta \in [0, 2\pi)\}$. Then $G \cong S^1$, and $\text{Fix } G = \mathbb{R}$ and consists of the constant functions. Moreover E, S^1 satisfy (*).

Below B_r denotes the closed ball in E of radius r centered at the origin and F^\perp denotes the orthogonal complement of a subspace F .

Theorem 3.5: Let E, S^1 satisfy (*) and let $\varphi \in C(B_R \cap E_k, E)$ be G equivariant with $\varphi = \text{id}$ on $(E_0 \cap B_R) \cup (E_k \cap \partial B_R)$. Then for any $\rho < R$, and $j < k$,

$$\begin{aligned} \text{Ind}((B_R \cap E_k) \cap E_j^\perp) &> \\ (3.6) \quad &> \text{Ind}(\{x \in B_R \cap E_k | \varphi(x) \in \partial B_\rho \cap E_j^\perp\}) > k-j. \end{aligned}$$

Proof: Let P_m denote the orthogonal projector of E onto E_m . Then for $m > k$, $P_m \varphi \in C(B_R \cap E_k, E_m)$ and is an equivariant map. Moreover $P_m \varphi = \text{id}$ on $(E_0 \cap B_R) \cup (E_k \cap \partial B_R)$. Hence by Proposition 3.3,

$$(3.7) \quad \text{Ind}(\{x \in B_R \cap E_k | \varphi(x) \in \partial B_\rho \cap E_m \cap E_j^\perp\}) > k-j.$$

Let K_m denote the argument of Ind in (3.7) and let

$$K = \{x \in B_R \cap E_k | \varphi(x) \in \partial B_\rho \cap E_j^\perp\}.$$

Then K is compact and $K \cap E_0 = \emptyset$. Hence by 3° of Lemma 3.1, $\text{Ind } K < \infty$ and there exists a $\delta > 0$ such that $\text{Ind } K = \text{Ind } N_\delta(K)$. We claim $K_m \subset N_\delta(K)$

$N_\delta(K)$ for m large. If so by 1° of Lemma 3.1, $\text{Ind } K_m < \text{Ind}_\delta N(K) = \text{Ind } K$ and (3.7) and 1° of Lemma 3.1 imply (3.6). If our claim were false, for all large m , there is an $x_m \in K_m \setminus N_\delta(K)$. Since $B_R \cap E_k$ is compact, we can assume $x_m \rightarrow \hat{x} \in (B_R \cap E_k) \setminus \text{int } N_\delta(K)$ as $m \rightarrow \infty$. Since

$$\|\varphi(\hat{x}) - P_m \varphi(x_m)\| \leq \|\varphi(\hat{x}) - P_m \varphi(\hat{x})\| + \|P_m(\varphi(\hat{x}) - \varphi(x_m))\| \rightarrow 0$$

as $m \rightarrow \infty$, $\varphi(\hat{x}) \in \partial B_j \cap E_j^\perp$, i.e. $\hat{x} \in K$. But $\hat{x} \notin \text{int } K$, a contradiction.

The proof is complete.

Remark 3.8: The above sort of situation occurs in the study of second order Hamiltonian systems [6]. Next we will study a case in which the domain of φ is not necessarily compact. This occurs in dealing with general Hamiltonian systems.

Theorem 3.9: Let E, S^1 satisfy (*) with $E_0^\perp = E^+ \oplus E^-$, where $E^+ = \overline{UE_m^+}$ (resp. $E_m^- = \overline{UE_m^-}$) and E_m^+, E_m^- are $2m$ dimensional mutually orthogonal invariant subspaces of E . Suppose $F_\rho \equiv E_0 \oplus E_m^+ \oplus E_m^-$ and $\varphi \in C(B_R \cap F_k, E)$ with φ equivariant, $\varphi = \text{id}$ on $(E_0 \cap B_R) \cup (F_k \cap \partial B_R)$ and $P^-\varphi(x) = \psi(x)P^-x + T(x)$ where P^- is the orthogonal projector of E onto E^- , $\psi \in C(E, [1, \alpha])$ and $T \in C(E, E^-)$ is compact. Then for any $\rho < R$ and $j > k$,

$$(3.10) \quad \text{Ind}(\varphi(B_\rho \cap F_k) \cap \partial B_\rho \cap F_j^\perp) > k-j.$$

Proof: Let Q_m denote the orthogonal projector of E onto $E_0 \oplus E^+ \oplus E_m^-$. Then $Q_m\varphi$ satisfies the hypotheses of Theorem 3.5 with E_k, E_j and E being replaced by F_k, F_j and Q_m . Hence by (3.6)

$$(3.11) \quad \text{Ind}(\{x \in B_R \cap Q_m F_k \mid Q_m\varphi(x) \in \partial B_\rho \cap F_j^\perp\}) > k-j.$$

Let K_m denote the argument of Ind in (3.11) and let

$$K = \{x \in B_R \cap F_k \mid \varphi(x) \in \partial B_\rho \cap F_j^\perp\}$$

Then K is compact. Indeed if (x_j) is a sequence in K , $x_j \equiv x_j^0 + x_j^+ + x_j^- \in E_0 \oplus E^+ \oplus E^-$. Since (x_j) is bounded and $B_R \cap F_j$ is closed and convex, we can assume x_j converges weakly to $x \in B_R \cap F_j$. Since

$(x_j^0), (x_j^+)$ lie in finite dimensional subspaces of E , we can assume these sequences converge strongly. Moreover $P^-\varphi(x_j) = 0 = \psi(x_j)x_j^- + T(x_j)$ or

$$(3.12) \quad x_j^- = -\psi(x_j)^{-1}T(x_j)$$

with $1 < \psi(x_j) < \alpha$. Thus the boundedness of (x_j) , compactness of T , and

(3.12) allow us to assume (x_j^-) also converges strongly. Hence

$\varphi(x_j) \rightarrow \varphi(x) \in \partial B_\rho \cap F_j^\perp$. Since $K \cap E_0 = \emptyset$, $i(K) < \infty$ by 3° of Lemma 3.1.

Continuing as in the proof of Theorem 3.5 with the additional information given by the form of $P^{-1}\varphi$ shows $\text{Ind } K_m < \text{Ind } K$ so (3.11) implies (3.10).

Remark 3.13: Results in the spirit of Theorems 3.5 and 3.9 have been obtained by Benci [9] for mappings φ which are equivariant homeomorphisms of E onto E of the form linear + compact.

Next a critical point theorem will be proved in which Theorem 3.5 plays an important role. If $I : E \rightarrow \mathbb{R}$, we say I is a G -invariant functional if $I(gx) = I(x)$ for all $g \in G$, $x \in E$. For $I \in C^1(E, \mathbb{R})$, we say I satisfies the Palais-Smale condition (PS) if any sequence (x_m) along which I is uniformly bounded and $I'(x_m) \rightarrow 0$ is a precompact sequence. Here $I'(x)$ denotes the Frechet derivative of I at x .

Suppose I satisfies

(I₁) For all finite dimensional subspaces \tilde{E} of E , there is an $r(\tilde{E}) > 0$ such that $I(x) < 0$ for $x \in \tilde{E}$ and $\|x\| > r(\tilde{E})$.

Choosing $\tilde{E} = E_0$, we see (I₁) and the continuity of I imply that $\sup_{E_0} I < \infty$.

Theorem 3.14: Suppose that E, S^1 satisfy (*) and $I \in C^1(E, \mathbb{R})$ is a G -invariant functional which satisfies (PS) and (I₁). If further I satisfies

(I₂) There is an $m_0 \in \mathbb{N}$ and $\rho > 0$ such that

$$(3.14') \quad I|_{\partial B_\rho \cap E_{m_0}^\perp} > \max(0, \sup_{E_0} I),$$

then I possesses an unbounded sequence of critical values.

Remark 3.15: A somewhat less general result in a \mathbb{Z}_2 setting was proved in [7]. One cannot obtain Theorem 3.14 by merely restricting to a subgroup of S^1 of order two and e.g. appealing to [7]. Indeed a special free \mathbb{Z}_2 action

was required in [7] and furthermore due to the possible presence of finite isotopy subgroups, restricting to Σ_2 may produce a fixed point set. Other Σ_2 and S^1 analogues of Theorem 3.14 can be found in Benci [9], [10], and Bahri [11].

The proof of Theorem 3.14 will be accomplished in several steps. Let $R_m \equiv r(E_m)$ obtained via (I_1) . Set $D_m = B_{R_m} \cap E_m$. Let $G_m = \{h \in C(D_m, E) | h \text{ is equivariant and } h(x) = x \text{ on } (E_0 \cup \partial B_{R_m}) \cap D_m\}$. Define

$$\Gamma_j = \{\overline{h(D_m \setminus Y)} | m > j, h \in G_m, Y \text{ invariant, } \text{Ind}(Y) < m-j\}.$$

Classes of sets somewhat like the Γ_j were used in [5]. The sets Γ_j possesses the following properties

- Lemma 3.16:
- (i) $\Gamma_{j+1} \subset \Gamma_j$.
 - (ii) If $B \in \Gamma_j$ and Z is an invariant set with $\text{Ind}(Z) < s < j$, then $\overline{B \setminus Z} \in \Gamma_{j-s}$.
 - (iii) If $\chi \in C(E, E)$ is equivalent and $\chi = \text{id}$ on $(E_0 \cup \partial B_{R_m}) \cap D_m$ for all $m > j$, then $\chi : \Gamma_j \rightarrow \Gamma_j$, i.e.

$$B \in \Gamma_j \implies \overline{\chi(B)} \in \Gamma_j.$$

Proof: (i) is immediate by the definition of Γ_j . For (ii), let $B \in \Gamma_j$. Therefore $B = \overline{h(D_m \setminus Y)}$ with $h \in G_m$ and $\text{Ind } Y < m-j$. Since $\overline{B \setminus Z} = \overline{h(D_m \setminus Y \cup h^{-1}(Z))}$ with $\text{Ind}(Y \cup h^{-1}(Z)) < \text{Ind}(Y) + \text{Ind}(h^{-1}(Z)) < \text{Ind}(Y) + \text{Ind}(Z) = m-j+s = m-(j-s)$ via 2° and 1° of Lemma 3.1, $\overline{B \setminus Z} \in \Gamma_{j-s}$. Lastly (iii) follows since if $B \in \Gamma_j$, $B = \overline{h(D_m \setminus Y)}$, then $\chi \circ h \in G_m$ and $\overline{\chi(h(D_m \setminus Y))} = \overline{\chi \circ h(D_m \setminus Y)}$.

With the aid of these sets Γ_j , we define a sequence of minimax values:

$$(3.17) \quad c_j = \inf_{B \in \Gamma_j} \sup_{u \in B} I(u), \quad j \in \mathbb{N}$$

Since $\Gamma_{j+1} \subset \Gamma_j$, we have $c_{j+1} \geq c_j$.

Lemma 3.18: For $j > m_0 + 1$, $c_j > c_{m_0+1} > \sup_{E_0} I$.

Proof: Let $B \in \Gamma_{m_0+1}$. Therefore $\overline{B} = h(D_m \setminus Y)$ where $h \in G_m$, $m > m_0 + 1$ and $\text{Ind}(Y) \leq m - m_0 - 1$. Let $\rho < R_m$. By Theorem 3.5, if $X = \{x \in D_m \mid h(x) \in \partial B_\rho \cap E_{m_0}^\perp\}$, $\text{Ind } X > m - m_0$. Therefore $\text{Ind } \overline{X \setminus Y} > m - m_0 - (m - m_0 - 1) = 1$ and $\overline{X \setminus Y} \neq \emptyset$. Since $h(\overline{X \setminus Y}) \subset W \equiv \overline{h(X \setminus Y)} \cap \partial B_\rho \cap E_{m_0}^\perp$ by 2° of Lemma 3.1, $W \neq \emptyset$. Hence if $w \in B \cap W$

$$(3.19) \quad \sup_B I > I(w) > \inf_W I > \inf_{\partial B_\rho \cap E_{m_0}^\perp} I.$$

Since (3.19) holds for all $B \in \Gamma_{m_0+1}$,

$$(3.20) \quad c_{m_0+1} > \inf_{\partial B_\rho \cap E_{m_0}^\perp} I$$

and the lemma follows from (3.14')

To continue we need a variant of a standard "Deformation Theorem". Let

$$K_c = \{x \in E \mid I(x) = c \text{ and } I'(x) = 0\} \text{ and } A_c = \{x \in E \mid I(x) < c\}.$$

Lemma 3.21: Let $I \in C^1(E, \mathbb{R})$ be G -invariant and satisfy (PS). Then for any $c \in \mathbb{R}$, $\bar{\epsilon} > 0$, and invariant neighborhood 0 of K_c , there exists $\epsilon \in (0, \bar{\epsilon})$ and $\eta \in C([0, 1] \times E, E)$ such that:

- 1° $\eta(t, x) = x$ if $x \in I^{-1}([c - \bar{\epsilon}, c + \bar{\epsilon}])$
- 2° $\eta(1, A_{c+\epsilon} \setminus 0) \subset A_{c-\epsilon}$
- 3° If $K_c = \emptyset$, $\eta(1, A_{c+\epsilon}) \subset A_{c-\epsilon}$
- 4° $\eta(t, x)$ is G -equivariant for each $t \in [0, 1]$.

Proof: The result without the G equivariance or invariance statements can be found in [12] or [13]. To obtain 4° also we need only modify these proofs by averaging over G to obtain a G equivariant pseudogradient vector field as in [9] or [14].

Now we can show that the c_j 's are indeed critical values of I together with a multiplicity statement:

Lemma 3.22: For each $j > m_0 + 1$, c_j is a critical value of I . Moreover if $c_{j+1} = \dots = c_{j+p} \equiv c$, $\text{Ind}(K_c) > p$.

Proof: It suffices to prove the stronger multiplicity assertion. Thus suppose $\text{Ind}(K_c) < p-1$. BY (PS), K_c is compact. It is also invariant. Hence by 3° of Lemma 3.1, there is a $\delta > 0$ such that $i(N_\delta(K_c)) = i(K_c)$. By Lemma 3.21 with $0 = \text{int } N_\delta(K_c)$ and $\bar{\epsilon} = \frac{1}{2}(c_{m_0+1} - \max(\sup_{E_0} I, 0)) > 0$, there is an $\epsilon \in (0, \bar{\epsilon})$ and mapping η such that

$$(3.23) \quad \eta(1, A_{c+\epsilon} \setminus N_\delta(K_c)) \subset A_{c-\epsilon}.$$

Choose $B \in \Gamma_{j+p}$ such that

$$(3.24) \quad \sup_B I < c + \epsilon$$

By (ii) of Lemma 3.16, $\overline{B \setminus N_\delta(K_c)} \in \Gamma_{j+1}$. By our choice of $\bar{\epsilon}$, $\eta(1, \cdot) = \text{id}$ on $(E_0 \cup \partial B_{R_m}) \cap D_m$ for all $m \in \mathbb{N}$. Hence by (iii) of Lemma 3.16,

$$\overline{\eta(1, B \setminus N_\delta(K_c))} \equiv Q \in \Gamma_{j+1}. \text{ Hence}$$

$$(3.25) \quad \sup_Q I > c_{j+1}$$

while by (3.23)-(3.24),

$$(3.26) \quad \sup_Q I < c_{j+1} - \epsilon,$$

a contradiction.

Remark 3.27: Note that for $j > m_0 + 1$, $K_{c_j} \cap E_0 = \emptyset$ via Lemma 3.18. Hence our multiplicity statement is not due to any contribution from $\text{Fix } G$.

To complete the proof of Theorem 3.14, we will show

Lemma 3.28: $c_j \rightarrow \infty$ as $j \rightarrow \infty$.

Proof: Since $c_{j+1} > c_j$, if $c_j \neq \infty$ as $j \rightarrow \infty$, $c_j \rightarrow \bar{c} < \infty$. Arguing as in [7], let $K = \{x \in E \mid c_{m_0+1} < I(x) < \bar{c} \text{ and } I'(x) = 0\}$. By (PS) K is compact and by Remark 3.27 and 3° of Lemma 3.1 $\text{Ind}(K) < \infty$ and for some $\delta > 0$, $\text{Ind}(N_\delta(K)) = \text{Ind } K$. Let $j = \text{Ind } K$. By Lemma 3.21 with $c = \bar{c}$ and $\bar{c} = \bar{c} - c_{m_0+1}$, there is an $\varepsilon \in (0, \bar{c})$ and equivariant $\eta(1, \cdot): A_{\bar{c}+\varepsilon} \setminus \text{int } N_\delta(K) \rightarrow A_{\bar{c}-\varepsilon}$. Let m be the smallest integer $> m_0+1$ such that $c_m > \bar{c} - \varepsilon$. Let $B \in \Gamma_{m+j}$ such that

$$\sup_B I < \bar{c} + \varepsilon$$

As in Lemma 3.22, $\overline{B \setminus N_\delta(K)}$ and $\overline{\eta(1, B \setminus N_\delta(K))} \equiv Q$ belong to Γ_m . Consequently

$$c_m < \max_Q I < \bar{c} - \varepsilon < c_m,$$

a contradiction.

Remark 3.29: In applications [6], one generally has $I(x) = \frac{1}{2}Q(x) + b(x)$ where Q is a quadratic form with Q positive definite on E_k^1 for some $k > 0$ and b is weakly continuous with $b(0) = 0$. Hence for any fixed

$$\rho > 0, \quad Q|_{\partial_{B_\rho} \cap E_m^1} > \beta_m \rho^2 \quad \text{for } m > k \text{ and } \beta_m \text{ bounded away from 0 while}$$

$$b|_{\partial_{B_\rho} \cap E_m^1} \rightarrow 0 \text{ as } m \rightarrow \infty \text{ since } x_m \in \partial_{B_\rho} \cap E_m^1 \text{ implies } x_m \text{ converges weakly}$$

to 0 and therefore $b(x_m) \rightarrow 0$ by the weak continuity of b . Thus (3.15) is satisfied. Hypotheses (I_1) is satisfied if b is "superquadratic" i.e. grows more rapidly than quadratically in an appropriate sense. See [7].

Remark 3.30: The novelty of Theorem 3.14 is not so much the conclusion of the theorem, since it is close to results of [9] and [11] but in the minimax characterization it provides for the critical values c_j . This characterization has proved to be useful in some recent perturbation results [6].

REFERENCES

1. Schwartz, J. T., Nonlinear Functional Analysis, Gordon and Breach, New York, 1969.
2. Granas, A., The theory of compact vector fields and some of its applications to the topology of function spaces, I, Rozprawy Matematyczne 30, Warsaw, 1962.
3. Holm, P. and E. H. Spanier, Involutions and Fredholm maps, Topology, 10, 1971, 203-218.
4. Rabinowitz, P. H., A note on a nonlinear elliptic equation, Indiana Univ. Math. J., 22, 1972, 43-49.
5. Fadell, E. R. and P. H. Rabinowitz, Generalized cohomological index theories for Lie group actions with an application to bifurcation questions for Hamiltonian systems, Inv. math., 45, 1978, 139-174.
6. Rabinowitz, P. H., Multiple critical points of perturbed symmetric functionals, to appear Trans. Amer. Math. Soc.
7. Ambrosetti, A. and P. H. Rabinowitz, Dual variational methods in critical point theory and applications, J. Functional Analy., 14, 1973, 349-381.
8. Spanier, E., Algebraic Topology, McGraw-Hill, New York, 1966.
9. Benci, V., On critical point theory for indefinite functionals in the presence of symmetries, to appear Trans. Amer. Math. Soc.
10. Benci, V., A geometrical index for the group S^1 and some applications to the research of periodic solutions of O.D.E.'s. To appear Comm. Pur. Appl. Math.
11. Bahri, A., Une methode perturbative en theorie de Morse, preprint.

12. Clark, D. C., A variant of the Ljusternik-Schnirelman theory, Indiana Univ. Math. J., 22, 1972, 65-74.
13. Rabinowitz, P. H., Variational methods for nonlinear eigenvalue problems, Proc. Sym. on Eigenvalues of Nonlinear Problems, Edizioni Cremonese, Rome, 1974, 141-195.
14. Ekeland, I., J. M. Lasry, On the number of periodic trajectories for a Hamiltonian flow on a convex energy surface, Ann of Math.
15. Bredon, G. E., Introduction to Compact Transformation Groups, Academic Press 1972, London, New York.

ERF:SYH:PHR/db

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2301	2. GOVT ACCESSION NO. AD-A140	3. RECIPIENT'S CATALOG NUMBER 458
4. TITLE (and Subtitle) BORSUK-ULAM THEOREMS FOR ARBITRARY S^1 ACTIONS AND APPLICATIONS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) E. R. Fadell, S. Y. Husseini, and P. H. Rabinowitz		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041 MCS-8110556
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18		12. REPORT DATE November 1981
		13. NUMBER OF PAGES 26
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Equivariant cohomology, index theory, Borsuk-Ulam theorem, intersection theorems, minimax, critical point, Hamiltonian system		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An S^1 version of the Borsuk-Ulam Theorem is proved for a situation where $\text{Fix } S^1$ may be nontrivial. The proof is accomplished with the aid of a new relative index theory. Applications are given to intersection theorems and the existence of multiple critical points is established for a class of functionals invariant under an S^1 symmetry.		

END

DATE
FILMED

3-82

DTIC